



Javanbakht T. (2022). Automated decision-making with TOPSIS for water analysis. *Journal of Engineering Sciences*, Vol. 9(1), pp. H19-H24, doi: 10.21272/jes.2022.9(1).h3

Automated Decision-Making with TOPSIS for Water Analysis

Javanbakht T.

Department of Computer Science, University of Quebec in Montreal,
201, President Kennedy St., Montreal, Quebec H2X 3Y7, Canada

Article info:

Submitted: April 4, 2022
Accepted for publication: June 3, 2022
Available online: June 6, 2022

*Corresponding email:

javanbakht.taraneh@courrier.uqam.ca

Abstract. This paper aims to present a new application of TOPSIS with an automated decision-making process for the analysis of drinking water. For this purpose, the algorithm was modified with a fuzzy disjunction, and the maximal output values were set to one. The properties of drinking water, such as total dissolved solids, hardness, electrical conductivity, and cost, were the criteria analyzed in this study. These criteria were analyzed with unmodified and modified algorithms. Therefore, the modified TOPSIS was also used to optimize the parameters of the candidates. The appearance of the value of 1.0 in the algorithm's output was due to the confusion of an individual's categories of drinking water and undrinkable water. The advantage of this investigation was that, for the first time, it allowed automated decision-making to detect the drinking water in different samples and analyze them according to their characteristics. This would be important in developing new technologies for detecting and analyzing drinking water in the environment. The results of this paper can be applied in materials sciences and engineering.

Keywords: TOPSIS, water, automated decision-making, computational engineering, process innovation.

1 Introduction

Drinking water is essential for human life and is in high demand in many countries [1-3]. Many people worldwide do not have access to safe drinking water [4]. The chemistry and toxicology of the chemicals in underground water can determine their characteristics for performing essential drinking water production processes [5].

The Technique for Order of Preference by Similarity to Ideal Solution (TOPSIS) is an appropriate decision-making method with diverse applications in science and engineering [6-10]. The candidates' ranking in this method is performed according to their distances from ideal solutions and their closeness coefficients [11-15].

The characteristics of drinking water, such as its total dissolved solids, hardness, electrical conductivity, and cost, must be analyzed [16-19]. The modified version of the TOPSIS method can help determine which water sample would be more appropriate according to its rank.

Moreover, the automated decision-making process must be performed with the modified algorithm. This would make the distinction between drinking water and non-consumable water. This analysis would help the next generation of robots distinguish and analyze different drinking water samples without human intervention. As

the amount of drinking water can decrease during the next decades on the planet, researching this material on other planets would be essential for human survival.

Demarcator theory is a theory according to which three characteristics, called the demarcators, are found in the category members for their distinction from the members of other categories. The strong and weak demarcators have an impact on this distinction.

However, the marginal demarcators do not positively affect this distinction, but their consideration leads to category confusion and inconsistency in epistemic beliefs [8, 20, 21].

According to the demarcator theory, the total dissolved solids in samples is their strong demarcator that impacts the distinction of drinking water from undrinkable water. In contrast, hardness and electrical conductivity are the weak demarcators that have less effect for this distinction. The transparency and amount of water and its cost cannot affect the importance of drinking water and undrinkable water. These marginal demarcators do not help humans distinguish the members of these categories.

This work aimed to optimize the drinking and undrinkable water samples with unmodified and modified TOPSIS methods.

For this purpose, the following tasks were performed:

1) optimization of water samples with unmodified TOPSIS;

2) modification of TOPSIS with the Łukasiewicz fuzzy disjunction;

3) optimization of water samples with modified TOPSIS.

The prediction, detection, and analysis of drinking water samples with modified TOPSIS with an automated decision-making process have not been performed previously. The results of this paper can find their applications in science and engineering.

2 Research Methodology

2.1 TOPSIS method

The TOPSIS code in Python is available on the GitHub website <https://github.com/Glitchfix/TOPSIS-Python/blob/master/topsis.py>. It was used for the optimization of parameters in this paper.

The main steps of this code were described previously [12].

2.2 Modified TOPSIS

The modified TOPSIS, including the Łukasiewicz fuzzy disjunction developed in new software, was used in this paper as described previously [8]. In the data analysis with the modified algorithm, the members of two categories of drinking water (candidates C1 and C2) and undrinkable water (C3, C4, and C5) were considered.

The category confusion due to humans' inappropriate consideration of the criteria that led to the inconsistency of their epistemic beliefs was analyzed as explained previously [8]. In the second series of analyses with the modified algorithm, turbidity as a marginal demarcator of water, which was a cost criterion, was added to the evaluation matrix. The categories confusion due to the consideration of this criterion as a profit criterion was evaluated with the fuzzy disjunction. This was because individuals mistakenly considered it a characteristic that positively distinguished the two water categories. The maximal value of 1.0, according to the Łukasiewicz fuzzy disjunction was observed in the evaluation matrix in the output.

3 Results and discussion

The first series of results were obtained with the unmodified TOPSIS algorithm.

Table 1 shows the evaluation matrix of water samples as candidates and their criteria.

Table 1 – Evaluation matrix of water samples as candidates and their criteria

Candidates/ Criteria	Total dissolved solids	Hardness	Electrical conductivity	Cost
C ₁	very low	very low	low	high
C ₂	very low	very low	low	medium
C ₃	medium	medium	medium	low
C ₄	medium	medium	medium	medium
C ₅	high	high	high	medium

Tables 2 and 3 show the matrices of triangular fuzzy data and their mean values, respectively.

Tables 4 and 5 show the matrix of the weights applied for each criterion of the water samples and the criteria matrix, respectively.

Table 6 shows the distances between the best and worst alternatives, the similarity coefficients, and rankings for water samples.

Table 2 – Matrix of triangular fuzzy data

Candidates/ Criteria	Total dissolved solids	Hardness	Electrical conductivity	Cost
C ₁	0.1, 0.2, 0.3	0.1, 0.2, 0.3	0.2, 0.3, 0.4	0.7, 0.8, 0.9
C ₂	0.1, 0.2, 0.3	0.1, 0.2, 0.3	0.2, 0.3, 0.4	0.4, 0.5, 0.6
C ₃	0.4, 0.5, 0.6	0.4, 0.5, 0.6	0.4, 0.5, 0.6	0.2, 0.3, 0.4
C ₄	0.4, 0.5, 0.6	0.4, 0.5, 0.6	0.4, 0.5, 0.6	0.4, 0.5, 0.6
C ₅	0.7, 0.8, 0.9	0.7, 0.8, 0.9	0.7, 0.8, 0.9	0.4, 0.5, 0.6

Table 3 – Matrix of the mean values of triangular fuzzy data

Candidates/ Criteria	Total dissolved solids	Hardness	Electrical conductivity	Cost
C ₁	0.2	0.2	0.3	0.8
C ₂	0.2	0.2	0.3	0.5
C ₃	0.5	0.5	0.5	0.3
C ₄	0.5	0.5	0.5	0.5
C ₅	0.8	0.8	0.8	0.5

Table 4 – Weights applied for each criterion of water samples

Alternatives/ Values	Total dissolved solids	Hardness	Electrical conductivity	Cost
C ₁ – C ₅	0.5	0.5	0.5	0.5

Table 5 – Criteria matrix for water samples

Alternatives/ Values	Total dissolved solids	Hardness	Electrical conductivity	Cost
C ₁ – C ₅	false	false	false	false

Table 6 – The distances from the best and worst alternatives (d_i^* and d_i^-), the similarity coefficients (CC_i) and rankings for water samples

Candidates	d_i^*	d_i^-	CC_i	Ranking
C ₁	0.1027	0.2207	0.6824	2
C ₂	0.0411	0.2292	0.8479	1
C ₃	0.1054	0.1550	0.5952	3
C ₄	0.1132	0.1315	0.5374	4
C ₅	0.2245	0.0617	0.2154	5

The second series of results were obtained with the modified TOPSIS algorithm. The Łukasiewicz fuzzy disjunction was applied to allow only the values equal to or below 1.0 to appear in the output.

Table 7 shows the evaluation matrix of water samples as candidates and their criteria.

Tables 8 and 9 show the matrices of triangular fuzzy data and their mean values, respectively.

Table 7 – Evaluation matrix of water samples as candidates and their criteria

Candidates/ Criteria	Total dissolved solids	Hardness	Electrical conductivity	Cost	Turbidity
C ₁	very low	very low	low	high	very low
C ₂	very low	very low	low	medium	very low
C ₃	medium	medium	medium	low	medium
C ₄	medium	medium	medium	medium	medium
C ₅	high	high	high	medium	medium

Table 8 – Matrix of triangular fuzzy data

Candidates/ Criteria	Total dissolved solids	Hardness	Electrical conductivity	Cost	Turbidity
C ₁	0.1,0.2,0.3	0.1,0.2,0.3	0.2,0.3,0.4	0.7,0.8,0.9	0.1,0.2,0.3
C ₂	0.1,0.2,0.3	0.1,0.2,0.3	0.2,0.3,0.4	0.4,0.5,0.6	0.1,0.2,0.3
C ₃	0.4,0.5,0.6	0.4,0.5,0.6	0.4,0.5,0.6	0.2,0.3,0.4	0.4,0.5,0.6
C ₄	0.4,0.5,0.6	0.4,0.5,0.6	0.4,0.5,0.6	0.4,0.5,0.6	0.4,0.5,0.6
C ₅	0.7,0.8,0.9	0.7,0.8,0.9	0.7,0.8,0.9	0.4,0.5,0.6	0.4,0.5,0.6

Table 9 – Matrix of the mean values of triangular fuzzy data

Candidates/Criteria	Total dissolved solids	Hardness	Electrical conductivity	Cost	Turbidity
C ₁	0.2	0.2	0.3	0.8	0.2
C ₂	0.2	0.2	0.3	0.5	0.2
C ₃	0.5	0.5	0.5	0.3	0.5
C ₄	0.5	0.5	0.5	0.5	0.5
C ₅	0.8	0.8	0.8	0.5	1.0

In Table 9, in comparison with table 3, the maximal value of 1.0 was observed for the last alternative due to the Łukasiewicz fuzzy disjunction, which showed the confusion of both drinking categories and undrinkable water samples by an individual.

Tables 10 and 11 show the matrix of the weights applied for each criterion of the water samples and the criteria matrix, respectively.

Table 12 shows the distances between the best and worst alternatives, the similarity coefficients, and rankings for water samples.

Table 10 – Weights applied for each criterion of water samples

Alternatives/ Values	Total dissolved solids	Hardness	Electrical conductivity	Cost	Turbidity
C ₁ – C ₅	0.5	0.5	0.5	0.5	0.5

Table 11 – Criteria matrix for water samples

Alternatives/Values	Total dissolved solids	Hardness	Electrical conductivity	Cost	Turbidity
C ₁ – C ₅	false	false	false	false	true

Table 12 – The distances from the best and worst alternatives (d_i^+ and d_i^-), the similarity coefficients (CC_i), and rankings for water samples

Candidates	d_i^+	d_i^-	CC_i	Ranking
C ₁	0.1515	0.1766	0.5382	2
C ₂	0.1315	0.1833	0.5824	1
C ₃	0.1159	0.1329	0.5341	3
C ₄	0.1205	0.1155	0.4894	4
C ₅	0.1796	0.1365	0.4318	5

The comparison of the obtained results revealed several important issues. The drinking water samples, candidates 1 and 2, were ranked in the first two positions, whereas undrinkable water samples were ranked in the three last positions.

Moreover, the cost impacts the ranking of drinking water samples, and the candidate with a lower cost, the second candidate, was ranked in the first position. The candidates' distances from the best and worst alternatives and their similarity coefficients were different with the unmodified and modified algorithms.

However, the same rankings were obtained in both analyses. The appearance of the value of 1.0 in the algorithm's output was due to the confusion of an individual's categories of drinking water and undrinkable water. This could be attributed to his inconsistency in epistemic beliefs.

Previously, the characteristics of several materials were investigated [22-26]. Moreover, new materials have been explored for their diverse applications in sciences and engineering [27-30]. It has been shown that nanoparticles [31, 32], polymers [33-36], and nanocomposites [37] could be used for water treatment. TOPSIS has been used for the selection of polymers [38-41], nanomaterials [42-44], and machine process parameters [45]. It has been shown to be efficient for optimizing analytic procedures and normalization methods [46, 47], and materials [48-52].

Although TOPSIS has been used to optimize some materials, its modified version with the Łukasiewicz fuzzy disjunction has not previously been used for their optimization. In other words, none of these previous studies has been done with the automated decision-making process with modified TOPSIS. To optimize these materials, it would be required to perform the automated decision-making process with the modified TOPSIS as presented in this paper.

More investigations are needed to improve the analysis method of drinking and undrinkable water samples with the TOPSIS algorithm. In the next step, the water samples with different concentrations of ions will be investigated.

4 Conclusions

The article aims to explain the analysis results of drinking and undrinkable water samples with an automated decision-making process. The tasks of this study were performed with unmodified and modified TOPSIS algorithms. The modified TOPSIS method was obtained by adding the Łukasiewicz fuzzy disjunction to the algorithm. For this purpose, the maximal values of the membership degrees were set to one.

To perform the analysis, the modified TOPSIS algorithm with this fuzzy disjunction was used, giving comparable results to the unmodified algorithm.

Although the distances from the best and worst alternatives and similarity coefficients differed with the unmodified and modified algorithms, the same rankings were obtained in both analyses.

The appearance of the value of 1.0 in the output of the algorithm, due to the confusion of the categories of drinking water and undrinkable water by the individual, could be attributed to his inconsistency in epistemic beliefs.

Overall, the research presents a new application of TOPSIS for predicting and detecting water samples that could be applied in sciences and engineering.

References

1. Yadi, M.T. Determinants of demand for the packaged drinking water, *International Journal of Innovative Research in Engineering & Multidisciplinary Physical Sciences*, 10(4):53-60, 2022.
2. Zhu, R., Fang, Y. Application of a water supply-demand balance model to set priorities for improvements in water supply systems: A case study from the Koshi river basin, Nepal, *Int J Environ Res Public Health*, 19(3), 1606, 2022. <https://doi.org/10.3390/ijerph19031606>.
3. Threats to sources of drinking water and aquatic ecosystem health in Canada, National Water Research Institute, Environment Canada, NWRI Scientific Assessment Report Serie 1, 2001.
4. Clerico, E.A. The future of water reuse in America in *Research to Improve Water-use Efficiency and Conservation : Technologies and Practices*, 30-51, 2007.
5. Emanuel, E., Simon, Y., Joseph, O. Characterization of hardness in the groundwater of Port-Au-Prince, An overview on the health significance of magnesium in the drinking water, *Aqua-LAC*, 5(2) :35-43, 2013.
6. Chen, Y., Li, K.W, Liu, S.F. An OWA-TOPSIS method for multiple criteria decision analysis, An OWA-TOPSIS method for multiple criteria decision analysis, *Expert Systems with Applications*, 38(5):5205-5211, 2011. <https://doi.org/10.1016/j.eswa.2010.10.039>.
7. Huang, W., Huang, Y.Y. Research on the performance evaluation of Chongqing electric power supply bureaus based on TOPSIS, *Energy Procedia*, 14:899-905, 2012. <https://doi.org/10.1016/j.egypro.2011.12.1030>.
8. Javanbakht, T. *Modélisation et traitement informatique de l'inconsistance des croyances épistémiques*, Thesis, University of Quebec in Montreal, 2022.
9. Ozturk, D., Batuk, F. Technique for order preference by similarity to ideal solution (TOPSIS) for spacial decision problems, *Proceedings ISPRS*, 2011.
10. Rahim, R. et al. Technique for order of preference by similarity to ideal solution (TOPSIS) method for decision support system in top management, *Information and Communication Technology Business and Management*, 7(3):290-293, 2018.
11. Indahingwati, A., Wajdi, M.B.N., Susilo, D.E., Kurniasih, N., Rahim, R. Comparison analysis of TOPSIS and fuzzy logic methods on fertilizer selection, *International Journal of Engineering and Technology*, 7(2.3):109-114, 2018. <https://doi.org/10.14419/ijet.v7i2.3.12630>.
12. Javanbakht, T. Analysis of nanoparticles characteristics with TOPSIS for their manufacture optimization, *Journal of Engineering Sciences*, 9(2):C1-C8, 2022. [https://doi.org/10.21272/jes.2022.9\(2\).c1](https://doi.org/10.21272/jes.2022.9(2).c1).
13. Jumarni, R.F., Zamri, N. An integration of fuzzy TOPSIS and fuzzy logic for multi-criteria decision making problems, *International Journal of Engineering and Technology*, 7(2):102-106, 2018. <https://doi.org/10.14419/ijet.v7i2.15.11362>.
14. Alguliyev, R., Aliguliyev, R., Yusifov, F. Modified fuzzy TOPSIS + TFNs ranking model for candidate selection using the qualifying criteria, *Soft Computing*, 24(1), 2020. <https://doi.org/10.1007/s00500-019-04521-2>.
15. Varnamkhandi, M.J., Sadabadi, S.A., Venicheh, A.-H. A new index for TOPSIS based on relative distance to best and worst points, *International Journal of Information Technology and Decision Making*, 16(03):695-719, 2020. <https://doi.org/10.1142/S0219622020500145>.
16. Thirumalini, S., Joseph, K. Correlation between electrical conductivity and total dissolved solids in natural waters, *Malaysian Journal of Science*, 28(1):55-61, 2009. <https://doi.org/10.22452/mjs.vol28no1.7>.
17. Büker, O. et al. Investigations on the influence of total water hardness and pH value on the measurement accuracy of domestic cold water meters, *Water*, 13, 2701, 2021. <https://doi.org/10.3390/w13192701>.
18. Clark, R.M., Goddard, H. Cost and quality of water supply, *Journal of American Water Works Associations*, 69(1):13-15, 1977. <https://doi.org/10.1002/j.1551-8833.1977.tb02533.x>.

19. Xianhong, Y. et al. Application analysis of conductivity in drinking water quality analysis, *International Energy, Environment and Water Resources Conference*, IOP publishing, 784, 012028, 2021. <https://doi.org/10.1088/1755-1315/784/1/012028>.
20. Javanbakht, T. Logique floue et arborescence comme outils de modélisation des catégories en tant que prototypes, Thesis, University of Quebec in Montreal, 2016.
21. Javanbakht, T. Être et Pensée, J. P. Beaudin & S. Robert (Eds.), BouquinBec, Montreal, 2020.
22. Javanbakht T, Laurent S, Stanicki D, David E. (2019). Related physicochemical, rheological, and dielectric properties of nanocomposites of superparamagnetic iron oxide nanoparticles with polyethyleneglycol, *Journal of Applied Polymer Science*, 136, 48280–48290. <https://doi.org/10.1002/app.48280>.
23. Javanbakht, T., Sokolowski, W. Thiol-ene/acrylate systems for biomedical shape-memory polymers. *Shape Memory Polymers for Biomedical Applications*, 157-166, 2015. <https://doi.org/10.1016/B978-0-85709-698-2.00008-8>.
24. Vollath, D., Szabó, D.V. Synthesis and properties of nanocomposites, *Advanced Engineering Materials*, 6(3):117-127, 2004. <https://doi.org/10.1002/adem.200300568>.
25. Djavanbakht, T., Carrier, V., André, J. M., Barchewitz, R., Troussel, P. Effets d'un chauffage thermique sur les performances de miroirs multicouches Mo/Si, Mo/C et Ni/C pour le rayonnement X mou. *Journal de Physique IV*, France, 10, 281-287, 2000. <https://doi.org/10.1051/jp4:20001031>.
26. Javanbakht, T., Laurent, S., Stanicki, D., Frenette, M. Correlation between physicochemical properties of superparamagnetic iron oxide nanoparticles and their reactivity with hydrogen peroxide. *Canadian Journal of Chemistry*, 98, 601-608, 2020. <https://doi.org/10.1139/cjc-2020-0087>.
27. Kaur, M., Tripathi, P.K. The basic properties of graphene and its applications, *International Journal of Research and Analytical Reviews*, 937-944, 2018.
28. Javanbakht, T., Ghane-Motlagh, B., Sawan, M. Comparative study of antibiofilm activity and physicochemical properties of microelectrode arrays. *Microelectronic Engineering*, 229, 111305, 2020. <https://doi.org/10.1016/j.mee.2020.111305>.
29. Javanbakht, T., Hadian, H., Wilkinson, K. J. Comparative study of physicochemical properties and antibiofilm activity of graphene oxide nanoribbons. *Journal of Engineering Sciences*, 7(1), C1-C8, 2020. [https://doi.org/10.21272/jes.2020.7\(1\).c1](https://doi.org/10.21272/jes.2020.7(1).c1).
30. Javanbakht, T., David, E. Rheological and physical properties of a nanocomposite of graphene oxide nanoribbons with polyvinyl alcohol. *Journal of Thermoplastic Composite Materials*, 0892705720912767, 2020. <https://doi.org/10.1177/0892705720912767>.
31. Fiorati, A., Bellingeri, A., Punta, C., Corsi, I., Venditti, I. (2020). Silver nanoparticles for water pollution monitoring and treatments: Ecosafety challenge and cellulose-based hybrids solution. *Polymers*, Vol. 12(8), 1635. <https://doi.org/10.3390/polym12081635>.
32. Nagar, A., Pradeep, T. Clean water through nanotechnology: Needs, gaps, and fulfillment, *ACS Nano*, 14(6):6420–6435, 2020. <https://doi.org/10.1021/acsnano.9b01730>.
33. Giwa, A. et al. *Polymeric materials for clean water*, Ed. R. Das, Springer, 167-190, 2019.
34. Khodakarami, M., Bagheri, M. Recent advances in synthesis and application of polymer nanocomposites for water and wastewater treatment, *Journal of Cleaner Production*, 296, 126404, 2021. <https://doi.org/10.1016/j.jclepro.2021.126404>.
35. Serajuddin, Chowdhury, A.I. et al. Application of a polymer in drinking water treatment: A case study, *5th International Conference on Civil Engineering for Sustainable Development*, 2020.
36. Adeola, A.O., Nomngongo, P.N. Advanced polymeric nanocomposites for water treatment applications: A holistic perspective, *Polymers*, 14(12):2462, 2022. <https://doi.org/10.3390/polym14122462>.
37. Beyene, H.D., Ambaye, T.G. Application of sustainable nanocomposites for water purification process, in *Sustainable Polymer Composites and Nanocomposites*, pp.387-412, 2019. https://doi.org/10.1007/978-3-030-05399-4_14.
38. Ardhiyanto, N.K., Pujiyanto, E., Rosyidi, C.N. Multi responses optimization of plastic injection molding for biodegradable polymers using Taguchi method and TOPSIS, *AIP Conference Proceedings*, 2097(1):030064, 2019. <https://doi.org/10.1063/1.5098239>.
39. Alaaeddin, M.H. et al. Polymer matrix materials selection for short sugar palm composites using integrated multi criteria evaluation method, *Composites B: Engineering*, 176, 107342, 2019. <https://doi.org/10.1016/j.compositesb.2019.107342>.
40. Narayanan, N.S. et al. Evaluation and optimization of surface roughness and metal removal rate through RSM, GRA, and TOPSIS techniques in turning PTFE polymers, *Advances in Manufacturing Technology*, 595-605, 2019. https://doi.org/10.1007/978-981-13-6374-0_65.
41. Chohan, J.S. et al. Taguchi S/N and TOPSIS based optimization of fused deposition modelling and vapor finishing process for manufacturing of ABS plastic parts, *Materials*, 13(22): 5176, 2020. <https://doi.org/10.3390/ma13225176>.
42. Yadav, R., Lee, H.-H. Fabrication, characterization, and selection using FAHP-TOPSIS technique of zirconia, titanium oxide, and marble dust powder filled dental restorative composite materials, *Polymers Advanced Technologies*, 33(1):3286-3295, 2022. <https://doi.org/10.1002/pat.5780>.
43. Zhang, K., Zhan, J., Yao, Y. TOPSIS method based on a fuzzy covering approximation space: An application to biological nano-materials selection, *Information Sciences*, 502, 297-309. <https://doi.org/10.1016/j.ins.2019.06.043>.
44. Loganathan, T.M. et al. Effect of *Cyrtostachys renda* fiber loading on the mechanical, morphology, and flammability properties of multi-walled carbon nanotubes/phenolic bio-composites, *Nanomaterials*, 11(11):3049, 2021. <https://doi.org/10.3390/nano11113049>.

45. Shunmugesh, K., Panneerselvam, K. Optimization of machine process parameters in drilling of CFRP using multi-objective Taguchi technique, TOPSIS and RSA techniques, *Polymers and Polymer Composites*, 25(3):185-192, 2017.
46. Al-Hazmi, H. et al. Application of TOPSIS for selection and assessment of analytical procedures for ibuprofen determination in wastewater, *Current Analytical Chemistry*, 12(4): 261-267, 2016. <https://doi.org/10.2174/1573411012666151009194541>.
47. Yang, W.-C., et al. Materials selection method using TOPSIS with some popular normalization methods, *Engineering Research Express*, 3, 015020, 2021. <https://doi.org/10.1088/2631-8695/abd5a7>.
48. Marzouk, M., El-Razek, M.A. Selecting demolition waste materials disposal alternatives using fuzzy TOPSIS technique, *International Journal of Nature Computing Research*, 6(2):38-57, 2020. <https://doi.org/10.4018/IJNCR.2017070103>.
49. Chen, C.-H. A hybrid multi-criteria decision-making approach based on ANP-entropy TOPSIS for building materials supplier selection, *Entropy*, 23(12):1597. <https://doi.org/10.3390/e23121597>.
50. Banwet, D.K., Majumdar, A. Comparative analysis of AHP-TOPSIS and GA-TOPSIS methods for selection of raw materials in textile industries, *Proceedings of the 2014 International Conference on Industrial Engineering and Operations Management*, 2071-2080, 2014.
51. Rao, A.Y. Koonal, R. Selection of optimum hybrid composite material for structural applications through TOPSIS technique, *International Journal of Surface Engineering and Interdisciplinary Materials Science*, 10(1):1-15, 2022.
52. Tiwari, S.K., Pande, S. Selection of gear materials using MCDM-TOPSIS approach, *International Journal of Manufacturing and Materials Processing*, 3(2), 2017. <https://doi.org/10.37628/ijmmp.v3i2.411>.